

| | | | |
|---|---|---------------------------------------|--------------------------------|
| Fiscal Year: | FY 2017 | Task Last Updated: | FY 03/16/2017 |
| PI Name: | Dulchavsky, Scott A. M.D., Ph.D. | | |
| Project Title: | Fluid Shifts | | |
| Division Name: | Human Research | | |
| Program/Discipline: | HUMAN RESEARCH | | |
| Program/Discipline--Element/Subdiscipline: | HUMAN RESEARCH--Biomedical countermeasures | | |
| Joint Agency Name: | TechPort: | No | |
| Human Research Program Elements: | (1) HHC: Human Health Countermeasures | | |
| Human Research Program Risks: | (1) Cardiovascular: Risk of Cardiovascular Adaptations Contributing to Adverse Mission Performance and Health Outcomes (2) SANS: Risk of Spaceflight Associated Neuro-ocular Syndrome (SANS) | | |
| Space Biology Element: | None | | |
| Space Biology Cross-Element Discipline: | None | | |
| Space Biology Special Category: | None | | |
| PI Email: | sdulchal@hfhs.org | Fax: | FY 313 916 9445 |
| PI Organization Type: | PUBLIC SERVICE | Phone: | 313 916 9306 |
| Organization Name: | Henry Ford Health System | | |
| PI Address 1: | Surgery | | |
| PI Address 2: | 2799 W. Grand Boulevard, CFP-1 | | |
| PI Web Page: | | | |
| City: | Detroit | State: | MI |
| Zip Code: | 48202-2608 | Congressional District: | 13 |
| Comments: | | | |
| Project Type: | FLIGHT | Solicitation / Funding Source: | 2011 Crew Health NNJ11ZSA002NA |
| Start Date: | 05/16/2013 | End Date: | 09/30/2018 |
| No. of Post Docs: | 0 | No. of PhD Degrees: | 0 |
| No. of PhD Candidates: | 0 | No. of Master' Degrees: | 0 |
| No. of Master's Candidates: | 0 | No. of Bachelor's Degrees: | 0 |
| No. of Bachelor's Candidates: | 0 | Monitoring Center: | NASA JSC |
| Contact Monitor: | Allcorn, Aaron | Contact Phone: | 281.244.8402 |
| Contact Email: | aaron.j.allcorn@nasa.gov | | |
| Flight Program: | ISS | | |
| Flight Assignment: | | | |
| Key Personnel Changes/Previous PI: | none | | |
| COI Name (Institution): | Ebert, Douglas (Wyle Laboratories, Inc.) Garcia, Kathleen (Wyle Laboratories, Inc.) Sargsyan, Ashot (Wyle Laboratories, Inc.) | | |
| Grant/Contract No.: | NNX13AK30G | | |
| Performance Goal No.: | | | |
| Performance Goal Text: | | | |

Task Description:

Editor's Note (7/11/2013): NOTE THIS IS A CONTINUATION OF FUNDING FOR NNX13AB42G (Microgravity Associated Compartmental Equilibration (MACE)) WITH THE SAME PRINCIPAL INVESTIGATOR.

Fifty percent of American astronauts have developed ocular refraction change after long duration space flight on the International Space Station (ISS). Recent findings have also included structural changes of the eye (papilledema, globe flattening, choroidal folds) and the optic nerve (sheath dilatation, tortuosity, and kinking), as well as imaging signs and lumbar puncture data indicative of elevated intracranial pressure (ICP). While short duration space flight is also characterized by vision disturbances, these are generally transient and do not appear to have lasting impacts on the structure or function of the eye. Changes in vision, eye, and adnexa morphology, are hypothesized to be the result of space flight-induced cephalad fluid shifts and transiently elevated intracranial pressure. This hypothesis, however, has not been systematically tested. In earlier anecdotal publications, ICP elevation in long-duration space flight has been inferred but without association with structural or functional changes of the eye. Furthermore, while fluid shifts and compartmentalization during short-duration space flight (Space Shuttle missions) have been studied, the fluid distribution patterns and their effects on intracranial pressure or the structure and function of the sensory organs in the course of long-duration space flight are not well known.

Several ISS crewmembers have reported consistent worsening of nasal congestion and associated symptoms in late afternoon hours, necessitating topical and systemic decongestant use. Although several explanations have been entertained, food (salt) and water intake are likely to have provoked these symptoms through postprandial modification of fluid balance or increase in the circulating volume that manifests in the most susceptible individuals.

The purpose of the proposed work is to objectively characterize the changes in fluid distribution, including intra/extracellular and intra/extravascular fluid shifts, by applying advanced non-invasive assessment technologies before, during, and after long duration space flight. Additionally, we will examine the relationship between the type and magnitude of the fluid shift with any effects on eye morphology and vision disturbances, intraocular pressure (IOP), and measures of intracranial pressure. Further, we seek to determine whether the magnitude of fluid shifts during space flight, as well as the above effects of those shifts can be predicted based upon crewmember baseline data and responses to acute head-down tilt tests performed before launch. Finally, we propose to evaluate the effect of lower body negative pressure (LBNP) on the above parameters.

To our knowledge, this is the first attempt to systematically determine the impact of the fluid distribution in microgravity on a comprehensive set of structural and functional measures including, but not limited to, those related to intracranial pressure, vision, morphology of the eye and its adnexa, and the vascular systems of the head and neck, during and after long duration space flight. The study design and methodology are based on the extensive relevant experience of the Investigators, including many successful ground-based, space flight analogue, and space flight projects and investigations.

Primary Hypothesis

Prolonged microgravity-induced, headward volume, and pressure shifts promote elevation of intracranial pressure and result in alterations in crewmembers' vision.

Specific Aims

Specific Aim I: To characterize fluid distribution and compartmentalization before, during, and after long-duration space flight.

Hypothesis 1: Fluid distribution measured by dilution techniques will reflect a headward fluid shift and an intra- to extra-vascular fluid shift during space flight, returning to pre-flight condition after landing.

Hypothesis 2: Regional headward fluid shifts in-flight are documented by increased cephalad venous dimensions (jugular veins) and flow characteristics, skin and soft tissue thickness.

Hypothesis 3: Fluid re-distribution towards the eye (detected in choroid, retina, and optic nerve head using ultrasonography and optical coherence tomography), and in arteries supplying ocular vascular beds, contributes to vision alterations.

Hypothesis 4: Splanchnic venous congestion (detected by portal vein size) contributes to headward volume shift, but is not in communication with the veins of head and neck. Thus, there should be a different level of venous congestion in these two compartments.

Specific Aim II: To correlate in-flight alterations of eye structure, ocular vascular parameters, and vision with headward fluid shifts, vascular dimensions, and flow patterns.

Hypothesis 5: Space flight-induced fluid shifts will have an upregulating effect on ICP and will alter ocular refraction / visual acuity. These changes will vary in magnitude and respectively, in their resolution pattern after space flight.

Hypothesis 6: In-flight ICP-related measures, IOP (intraocular pressure), venous and arterial morphometric and flow characteristics, vascular resistance of ocular vascular beds, and optic nerve anatomy will trend towards normal-gravity levels temporarily during and residually after fluid sequestration (LBNP) interventions.

Specific Aim III: To determine systemic and ocular factors of individual susceptibility to the development of ICP elevation and/or vision alterations.

Hypothesis 7: Subjects with greater fluid shifts (as measured by the ultrasound method in Aim 1) during pre-flight testing will experience greater fluid shifts in-flight and will be more susceptible to flight-induced vision alterations.

Hypothesis 8: Subjects who are resistant to the reversal of in-flight symptoms and physiological status through the application of LBNP will be more susceptible to persistent flight-induced vision alterations.

Hypothesis 9: Propensity towards more severe changes in-flight and the more indolent postflight resolution pattern will correlate with a range of individual characteristics of the crewmembers, such as anatomical and microanatomical and physiological features and potentially, hitherto unsuspected factors.

NOTE: This study was merged with investigations from Dr. Alan Hargens (Fluid distribution before, during and after prolonged space flight) and Dr. Michael Stenger (Distribution of Body Fluids during Long Duration Space Flight and Subsequent Effects on Intraocular Pressure and Vision Disturbance) resulting in a comprehensive study titled "Fluid

Shifts Before, During and After Prolonged Space Flight and Their Association with Intracranial Pressure and Visual Impairment” (short title: Fluid Shifts).

Rationale for HRP Directed Research:

Research Impact/Earth Benefits:

Current means of measuring increased intracranial pressure require an invasive monitoring system with skilled medical personnel. The techniques outlined in this proposal, if verified, would provide a rapid, accurate, non-invasive, and scalable solution to measure increases in intracranial pressure for a number of critical medical conditions. These studies will also provide physiological insight to the mechanisms of fluid shifts and their relationship to intracranial pressure. This information could be relevant to terrestrial disorders of intracranial pressure such as idiopathic intracranial hypertension (IIH).

We have made significant progress over the past year in all dimensions of the Fluid Shifts (FS) experiment. To date the team has performed preflight baseline data collection on eleven crewmembers, completed all inflight sessions for 10 subjects with the exception of the final return minus 45 days (R-45) time points for the last three subjects. Eleven postflight tilt table sessions have been completed for the first six subjects, ranging from return plus 10 days (R+10) to R+180. Upright and 3 Tesla (3T) postflight MRI were also collected on these six subjects (ranging from R+1 to R+48). All training is now complete, and the last four Fluid Shifts subjects are currently on ISS.

The FS team has continued to collectively respond to changes in circumstances (inflight schedules, Russian travel limitations, hardware failures, etc.). The Cochlear and Cerebral Fluid Pressure (CCFP) Analyzer's hard drive failed in August 2016 which resulted in the loss of inflight CCFP and tympanometry data until a replacement unit was delivered on orbit in November 2016. In January 2017 one of the otoacoustic emission analyzer's probes became clogged with earwax so the backup probe is being used for continued data collection. Procedures were developed and successfully implemented in February 2017 to clean the primary probe. The Fluid Shifts team has also developed several alternate data collection plans based on changing crew schedules, most recently including the potential for an extended mission for the Expedition 51 Commander.

Supine and tilted 3T MRI data collections continued at the Victory Lakes facility (UTMB--University of Texas Medical Branch), taking advantage of the existing MRID (MED B) pulse sequences. Subjects continued to tolerate the procedure well and all early assessments of data quality gave satisfactory results. Since 2016, the team includes Dr. Larry Kramer (UTHSC-H) for general MRI advising and assistance with data analysis. Specifically, work is underway for CSF (cerebrospinal fluid) flow quantification in the Sylvian aqueduct and determination of pre- and postflight CSF production rates. Other MRI analysis methods continue to be refined to optimize data analysis in terms of quality and resources. Despite the decision by Medical Operations to use IV contrast during MED B MRI scans as the preferred version for venography, none of the Fluid Shifts subjects has received contrast injections and data collections are likely to continue with non-contrast techniques.

Since Russian and European Space Agency (ESA) subjects will “direct return” to their respective countries, immediate postflight testing in Houston will not be possible. We have been working to standardize the imaging procedure and protocol elements among UTMB, Research Center of Neurology in Moscow, Russia, and DLR/envihab in Cologne, Germany. Both Partner facilities are equipped with -15 degree foam wedges and have prior experience with tilted MRI. A test run of all sequences was performed at DLR in a healthy volunteer (February 2017), and subsequent evaluation of data was performed. Additional testing may be conducted in March to complete standardization. A similar effort is underway with the Russian facility, which has demonstrated the ability to perform the protocol with satisfactory results. Additional efforts will be made to optimize the data in the 2 remaining postflight sessions.

The “free-floating” use of the OCT device for inflight Chibis sessions continues to work well for crewmembers, resulting in similar exam times and data quality when compared to the traditional chinrest method.

Similar to other measures, ultrasound data is collected in the inflight baseline state (Columbus module) and again during Chibis (Russian Service Module), which allows us to contrast space normal to the lower negative pressure state induced by the Chibis device. Transcranial Doppler, optic nerve sheath diameter, central retinal and ophthalmic artery Doppler, among other parameters, are being analyzed to compare conditions and individual subject variation.

Task Progress:

Our team has continued to make advances in the analysis of otoacoustic emission (OAE) data over the past year. Dr. David Kemp continues to provide invaluable OAE expertise and has further automated his OAE processing software. Rozela Melgoza returned for a second summer audiology internship in 2016 and continued to be a major contributor for data processing and exploration of data nuances under Dr. Kemp's guidance. We have focused our analysis on transient evoked OAE (TEOAE) phase shifts, which have been highly systematic in response to posture change and lower body negative pressure during preflight testing. TEOAE phase shifts are consistent with expected intracranial pressure changes due to HDT, and appear to be consistent across multiple subjects. The team was fortunate to host a second audiology summer intern in 2016 (Tyler Caldwell). Tyler performed ground studies to define the effect of probe positioning, middle ear pressure, and head-down tilt on OAE stimulus and response signals, which resulted in two posters related to Fluid Shifts, one at the NASA Human Research Program (HRP) Investigators' Workshop 2017 and one at the Texas Academy of Audiology. A third abstract related to this summer internship work has been submitted to the American Academy of Audiology meeting which will be held in April 2017.

Our team attended the NASA HRP Investigators' Workshop in Galveston, TX in January 2017, presenting an overall project lecture for Fluid Shifts and participating in many Vision Impairment and Intracranial Pressure (VIIP)-related sessions and discussions. Our team also presented two otoacoustic emissions posters based on methods developed and data collected for the Fluid Shifts project.

Presentations (past year):

1) Caldwell T, Ebert DJ, Danielson R, Kemp D, Le Prell C. Assessing the Utility of Otoacoustic Emissions for Monitoring Intracranial Pressure in Microgravity through Analog Observations. Presented at the 17th Annual Texas Academy of Audiology Conference, Dallas, TX, October 20-22, 2016.

2) Caldwell T, Ebert D, Kemp D, Danielson R, Stenger M. Effects of Middle Ear Pressure and Intracranial Pressure on Transient-Evoked Otoacoustic Emissions. Human Research Program Investigators' Workshop, Galveston, TX, January 23-26, 2017.

| | |
|---|---|
| | <p>3) Kemp D, Melgoza R, Ebert D, Danielson R, Stenger M, Hargens A, Dulchavsky S. Otoacoustic Emissions in Fluid Shift Studies: Methodology and Confounding Factors. Human Research Program Investigators' Workshop, Galveston, TX, January 23-26, 2017.</p> <p>4) Melgoza R, Kemp D, Ebert D, Danielson R, Stenger M, Hargens A, Dulchavsky S. A Longitudinal Study of Transient Evoked Otoacoustic Emissions in Relation to Spaceflight (Fluid Shifts). Human Research Program Investigators' Workshop, Galveston, TX, January 23-26, 2017.</p> <p>5) Stenger MB, Hargens AR, Dulchavsky SA, Arbielle P, Danielson RW, Ebert DJ, Garcia KM, Johnston SL, Laurie SS, Lee SMC, Liu J, Macias B, Martin DS, Minkoff L, Ploutz-Snyder R, Ribeiro LC, Sargsyan A, Smith SM. Fluid Shifts. Human Research Program Investigators' Workshop, Galveston, TX, January 23-26, 2017.</p> |
| Bibliography Type: | Description: (Last Updated: 02/23/2023) |
| Abstracts for Journals and Proceedings | <p>Caldwell T, Ebert DJ, Danielson R, Kemp D, Le Prell C. "Assessing the Utility of Otoacoustic Emissions for Monitoring Intracranial Pressure in Microgravity through Analog Observations." Presented at the 17th Annual Texas Academy of Audiology Conference, Dallas, TX, October 20-22, 2016.</p> <p>7th Annual Texas Academy of Audiology Conference, Dallas, TX, October 20-22, 2016. , Oct-2016</p> |
| Abstracts for Journals and Proceedings | <p>Caldwell T, Ebert D, Kemp D, Danielson R, Stenger M. "Effects of Middle Ear Pressure and Intracranial Pressure on Transient-Evoked Otoacoustic Emissions." 2017 NASA Human Research Program Investigators' Workshop, Galveston, TX, January 23-26, 2017.</p> <p>2017 NASA Human Research Program Investigators' Workshop, Galveston, TX, January 23-26, 2017. , Jan-2017</p> |
| Abstracts for Journals and Proceedings | <p>Kemp D, Melgoza R, Ebert D, Danielson R, Stenger M, Hargens A, Dulchavsky S. "Otoacoustic Emissions in Fluid Shift Studies: Methodology and Confounding Factor." 2017 NASA Human Research Program Investigators' Workshop, Galveston, TX, January 23-26, 2017.</p> <p>2017 NASA Human Research Program Investigators' Workshop, Galveston, TX, January 23-26, 2017. , Jan-2017</p> |
| Abstracts for Journals and Proceedings | <p>Melgoza R, Kemp D, Ebert D, Danielson R, Stenger M, Hargens A, Dulchavsky S. "A Longitudinal Study of Transient Evoked Otoacoustic Emissions in Relation to Spaceflight (Fluid Shifts)." 2017 NASA Human Research Program Investigators' Workshop, Galveston, TX, January 23-26, 2017.</p> <p>2017 NASA Human Research Program Investigators' Workshop, Galveston, TX, January 23-26, 2017. , Jan-2017</p> |
| Abstracts for Journals and Proceedings | <p>Stenger MB, Hargens AR, Dulchavsky SA, Arbielle P, Danielson RW, Ebert DJ, Garcia KM, Johnston SL, Laurie SS, Lee SMC, Liu J, Macias B, Martin DS, Minkoff L, Ploutz-Snyder R, Ribeiro LC, Sargsyan A, Smith SM. "Fluid Shifts." 2017 NASA Human Research Program Investigators' Workshop, Galveston, TX, January 23-26, 2017.</p> <p>2017 NASA Human Research Program Investigators' Workshop, Galveston, TX, January 23-26, 2017. , Jan-2017</p> |