

Fiscal Year:	FY 2019	Task Last Updated:	FY 12/06/2018
PI Name:	Porada, Christopher Ph.D.		
Project Title:	Effects of Microgravity on the Risks of Space Radiation-induced Leukemogenesis		
Division Name:	Space Biology		
Program/Discipline:			
Program/Discipline--Element/Subdiscipline:			
Joint Agency Name:		TechPort:	No
Human Research Program Elements:	None		
Human Research Program Risks:	None		
Space Biology Element:	(1) Cell & Molecular Biology (2) Animal Biology: Vertebrate		
Space Biology Cross-Element Discipline:	(1) Immunology		
Space Biology Special Category:	(1) Translational (Countermeasure) Potential		
PI Email:	cporada@wakehealth.edu	Fax:	FY
PI Organization Type:	UNIVERSITY	Phone:	336-713-1655
Organization Name:	Wake Forest University School of Medicine		
PI Address 1:	Regenerative Medicine		
PI Address 2:	391 Technology Way, 4th Floor		
PI Web Page:			
City:	Winston Salem	State:	NC
Zip Code:	27101-3074	Congressional District:	5
Comments:			
Project Type:	GeneLab,GROUND	Solicitation:	2016 Space Biology (ROSBio) NNH16ZTT001N-GeneLab. Appendix A: Translational Systems Biology and Informatics Research Using the GeneLab Data System
Start Date:	02/01/2017	End Date:	01/31/2020
No. of Post Docs:	1	No. of PhD Degrees:	
No. of PhD Candidates:	1	No. of Master' Degrees:	1
No. of Master's Candidates:		No. of Bachelor's Degrees:	
No. of Bachelor's Candidates:	2	Monitoring Center:	NASA ARC
Contact Monitor:	Sato, Kevin	Contact Phone:	650-604-1104
Contact Email:	kevin.y.sato@nasa.gov		
Flight Program:			
Flight Assignment:	NOTE: Extended to 1/31/2020 per F. Hernandez/ARC (Ed., 7/23/19) NOTE: Extended to 7/31/2019 per F. Hernandez/ARC (Ed., 2/14/19)		
Key Personnel Changes/Previous PI:	November 2017 report: No changes.		
COI Name (Institution):	Almeida-Porada, Maria Graca M.D., Ph.D. (Wake Forest University) Walker, Steve Ph.D. (Wake Forest University) Wilson, Paul Ph.D. (University of California, Davis)		
Grant/Contract No.:	NNX17AE49G		
Performance Goal No.:			
Performance Goal Text:			

Task Description:

We will specifically be making use of data generated as part of GeneLab experiment sets GLDS-53, GLDS-55, and GLDS-25 as the basis for the novel hypothesis to be tested in the current proposal: microgravity (μG) acts in concert with solar particle event (SPE) and galactic cosmic ray (GCR) radiation to produce deleterious effects on the human hematopoietic system, which may lead to an enhanced risk of leukemogenesis, as a result of both increased genomic damage to cells of the hematopoietic system, and a reduced ability of the immune system to recognize and clear hematopoietic cells that have undergone malignant transformation as a result of exposure to SPE/GCR radiation and conditions of microgravity. Data generated from the aforementioned GeneLab studies support this hypothesis, as these data have shown that μG : 1) induces higher levels of spontaneous DNA damage in human hematopoietic cells; 2) markedly alters the ability of mature human immune cells to respond appropriately to stimuli; 3) diminishes the ability of human lymphocytes to efficiently repair DNA damage in response to ionizing radiation; and 4) leads to alterations in the levels of multiple miRNAs that have been implicated in a variety of human hematopoietic malignancies. We have also generated a wealth of data to support the hypothesis that μG and space radiation likely act synergistically to increase astronaut risk of leukemogenesis during a prolonged mission beyond LEO (low Earth orbit). In the present proposal, we will build upon these data by performing studies to directly test the ability of μG to increase the risk of leukemic transformation in human hematopoietic stem/progenitor cells (HSC), while simultaneously reducing the ability of generated immune cells from recognizing and removing any malignant clones that arise.

Rationale for HRP Directed Research:**Research Impact/Earth Benefits:**

Our research has thus far revealed that conditions of microgravity lead to marked alterations in the ability of human hematopoietic stem/progenitor cells (HSC) to repair DNA double strand breaks (DSBs) that are characteristic of the damage that occurs following exposure to ionizing radiation. Moreover, microgravity also appears to impair the ability of human HSC to generate functional dendritic cells, which act as critical sentinels within the immune system, detecting infectious invaders and cells that have undergone malignant transformation, and alerting/priming immune effectors to eliminate these threats. Further adding to these deleterious effects, we have also found that microgravity negatively affects the ability of human natural killer (NK) cells to recognize and lyse human leukemic cells. Taken together, our results to date have shown that conditions of microgravity present during spaceflight could add to the risks of leukemogenesis as a result of exposure to space radiation, both by impairing the ability of human HSC to repair the induced damage and by crippling the generation and function of the immune cells needed to recognize and eliminate cells damaged by the radiation. An understanding of the mechanism(s) by which microgravity affects and impairs these different processes could lead to the development of novel methods to target and augment these pathways, and thereby enhance the processes of DNA repair and anti-tumor immunity. Such developments could have a profound impact on the treatment of cancer and on the lives of patients suffering from this disease.

One of the major factors limiting manned spaceflight beyond low-Earth orbit is the poorly defined health risks as a result of exposure to space radiation in the form of solar energetic particles (SEP) and galactic cosmic rays (GCR) that could potentially increase cancer morbidity/mortality in astronauts. We have recently demonstrated seriousness of this risk by showing that exposing human hematopoietic stem cells (HSC) to simulated GCR (at high dose-rate) results in the generation of leukemia when these cells repopulate the hematopoietic system of mice (Leukemia, 31(6), 1398-1407, 2017). In addition to being exposed to high energy radiation, astronauts are subjected to conditions of microgravity (μG) during spaceflight, which also exerts a wide range of untoward effects on the body, including altering immunity and the response to radiation. We therefore hypothesized that conditions of microgravity μG present during spaceflight may act in concert with SEP and GCR radiation to produce deleterious effects on the human hematopoietic system, leading to an enhanced risk of leukemogenesis, as a result of both increased genomic damage to cells of the hematopoietic system, and a reduced ability of the generated immune system to recognize and clear hematopoietic cells that have undergone malignant transformation.

In the 1st set of studies to test this hypothesis, we treated the human HSC-like KG1a cell line with an acute dose (10 $\mu\text{g/ml}$) of the radiomimetic drug bleomycin to mimic the damaging effects of SEP/GCR radiation. We then took the treated cells, in addition to an identical batch of untreated cells, and split them in half. Half of the cells from each aliquot were cultured under conditions of normal gravity (1G) in a 37 degrees C incubator, and the other half was cultured in the Synthecon Rotary Cell Culture System (RCCS) to create a state of continual freefall, and thereby mimic μG . At 1 and 4 hours, half of the cells were harvested from the 1G and μG cultures, and we quantitated the extent of double-strand breaks (DSBs) and the kinetics of repair, using flow cytometry and confocal imaging to monitor the formation and disappearance of gamma-H2AX foci. Interestingly, both the bleomycin-treated and untreated cells experienced a moderate increase in their median fluorescence intensity (MFI) when cultured in a μG environment, indicating that just the presence of conditions of simulated μG may lead to an increase in DNA damage.

To determine the impact conditions of μG exerted on the ability of KG1a cells to repair DSBs induced by the radiomimetic bleomycin, we calculated the ratio of bleomycin-treated cells' MFI to untreated cells' MFI under conditions of 1G vs. those in μG between the 1 hr and 4 hr time points. We reasoned that if DNA repair proceeds normally, this ratio should decrease from the 1 hr to 4 hr time point, since gamma-H2AX foci disappear as DNA is repaired. When we first examined the cells cultured in 1G, this ratio did indeed decrease, by 26%, from the 1 hr time point to the 4 hr time point for cells, demonstrating successful repair of the bleomycin-induced DNA damage during this time. In marked contrast, cells maintained in μG experienced a 20% increase over this same period, suggesting they were unable to repair the bleomycin-induced DNA damage, and this damage accumulated during the 4 hr incubation.

Since the "classical" method for assessing DSBs induced by bleomycin or ionizing radiation is to perform immunofluorescence microscopy on cells following staining with gamma-H2AX, we used this traditional method to confirm that fluorescence events that we were detecting on the flow cytometer were indeed due to the presence of gamma-H2AX foci within the nuclei of the human HSC-like KG1a cells maintained in conditions of simulated μG during the period of DNA repair following bleomycin exposure.

From the studies we have performed to-date, we conclude that HSC DNA damage repair is compromised in conditions of μG , leading to an accumulation of DSBs that cannot be resolved over time. These findings thus support our hypothesis that conditions of μG may enhance the genotoxic effects of space radiation, and increase the risk of leukemogenesis. These findings were recently published in Dr. Elizabeth Blaber's Special Issue of Stem Cells and Development that is focused on Microgravity (Stem Cells Dev. 2018 Sep 15;27(18):1257-1267. doi: 10.1089/scd.2018.0052).

Since last year's Progress Report, we have established a new collaboration with Dr. Pierre-Alexandre Vidi at the Wake Forest School of Medicine. Dr. Vidi owns a custom-made micro gamma-irradiator, that easily fits within a standard

tissue culture incubator. We have been working with Dr. Vidi for the past several months to find a way to adapt this device to enable us to irradiate human KG1a cells (and eventually primary human CD34+ HSC) while they are spinning in the RWV/HARV. We have now adapted the system to accomplish this task. We are thus poised, for the first time ever, to be able to study the incidence of ionizing radiation-induced DNA damage, and its subsequent repair, in human cells while they are in conditions of microgravity. Until this point, all studies examining the effects of microgravity on DNA damage and repair have been forced to remove the cells from microgravity during exposure to irradiation. We thus feel that these studies will provide highly novel and valuable data to NASA regarding this important issue.

Task Progress:

In our 2nd set of experiments, we tested the hypothesis that μG alters the development and functionality of critical regulators of the immune system, which could further enhance cancer risk during spaceflight. The generation of an effective immune response requires that antigens be processed and presented to T cells by antigen presenting cells (APC), the most potent of which are dendritic cells (DC), which stimulate naïve T cells and initiate primary immune responses. DC also function as effector cells in innate immunity. Because DC have influence over both the innate and acquired arms of immunity, a defect in their production and/or function would be expected to result in broad impairment of immunity. We therefore tested whether DC could be generated from human HSC in μG , again using the RCCS. CD34+ cells were cultured in 1G or μG using cytokine-supplemented serum-free media, and were analyzed for the presence of plasmacytoid (CD123+) and myeloid (CD11c+) DC by flow cytometry. While culture in both 1G and μG produced higher numbers of myeloid (CD11c+) vs. plasmacytoid (CD123+) DC, the HSC cultured in 1G differentiated into DC in significantly higher numbers than those cultured in μG . Moreover, culture in μG significantly delayed the generation of DC; while DC were generated within 7 days in 1G, they only appeared after 14 days in μG . Our results thus suggest that μG delays the production of DC and reduces the number of these important APC that are generated. A manuscript detailing these findings was also recently published in Dr. Elizabeth Blaber's Special Issue of Stem Cells and Development that is focused on Microgravity (Stem Cells Dev. 2018 Sep 15;27(18):1257-1267. doi: 10.1089/scd.2018.0052).

In our 3rd set of experiments, we have begun exploring the effects that conditions of μG exert on the generation and functionality of human NK cells, focusing on their ability to recognize and lyse human leukemic cells, as our prior Human Research Program (HRP)-supported work showed that exposure of human HSC to simulated GCR ions induced T-ALL. To begin these studies and optimize the methods/assays to be used, we began with the human IL-2-independent NK cell line NK-92MI, testing the effects that a brief (48 h) exposure to conditions of μG had on the ability of these cells to subsequently recognize and lyse the human erythroleukemic cell line K562, and the human T-ALL cell line MOLT-4. Even this relatively brief exposure to conditions of μG dramatically reduced the ability of this human NK cell line to recognize and lyse both of these human leukemic targets, irrespective of the effector:target (E:T) ratio used. After extensive optimization, we have finally managed to establish culture conditions that permit the long-term maintenance and expansion of primary human peripheral blood NK cells, and studies are now ongoing to assess whether conditions of μG exerts similar effects on the functionality of these primary cells. Samples have also been prepared to perform transmission electron microscopy (TEM) to define the subcellular alterations that occur in the NK cells exposed to μG that so dramatically affects their ability to recognize and/or lyse leukemic cells. Studies are focused on the alterations μG induces in the actin cytoskeleton of the NK cells, as the actin filaments are known to play a critical role in both the formation of the immune synapse and the movement and ultimate release of the cytolytic granules from the NK cells. In addition, we have joined together with several other faculty at Wake Forest Institute for Regenerative Medicine (WFIRM) to purchase a NanoString device, which will enable us to rapidly and thoroughly define all of the transcriptional changes (both mRNA and miRNA) that occur in both NK-92MI and primary human NK cells as a result of exposure to conditions of μG .

In addition to the afore-detailed studies in which NK cells were exposed to conditions of μG and then used for NK lysis assays to assess functionality, we reasoned that it would be far more interesting and biologically relevant to devise a means of assessing NK functionality while the NK cells were in conditions of μG . To accomplish this objective, we spent roughly 6 months adapting the nonradioactive NK lysis kit to enable us to be able to perform the whole assay in the RWV/HARV. We have now repeated all of our prior studies with the NK-92MI cell line, and we have shown that the effects of conditions of μG on NK cell activity against human leukemic cells are even more pronounced when the NK cells are allowed to find, engage, and lyse the leukemic cells while in conditions of μG . A manuscript detailing all of these important findings on human NK cells functionality in conditions of μG is currently being prepared for submission to Nature Microgravity.

Combining the data we have generated on the immune-altering effects of μG with those of the preceding experiments on DSB repair, we feel that our studies to-date support our overall hypothesis that conditions of μG present during spaceflight may enhance the risk of SEP/GCR-induced leukemogenesis, highlighting the importance of developing means of simulating gravity during spaceflight or identifying biological countermeasures to the deleterious effects of conditions of μG to reduce astronaut risk.

Bibliography Type:

Description: (Last Updated: 01/17/2019)

Abstracts for Journals and Proceedings

Kuhlman BM, Almeida-Porada G, Porada CD. "Microgravity Impairs Anti-Leukemic Activity of Human NK Cells." To be presented at the NASA Human Research Program Investigators' Workshop, Galveston, TX, 2019. 2019 NASA Human Research Program Investigators' Workshop, Galveston, TX, January 22-25, 2019. , Jan-2019

Abstracts for Journals and Proceedings

Kuhlman BM, Almeida-Porada G, Porada CD. "Microgravity Impairs Anti-Leukemic Activity of Human NK Cells." Poster presentation. 34th Annual Meeting of the American Society for Gravitational and Space Research, Bethesda, MD, October 31-November 3, 2018. 34th Annual Meeting of the American Society for Gravitational and Space Research, Bethesda, MD, October 31-November 3, 2018. , Nov-2018

Abstracts for Journals and Proceedings

Kuhlman BM, Almeida-Porada G, Porada CD. "Could Conditions of Microgravity Enhance Cancer Risk from Space Radiation?" Podium Talk. 6th Annual NextGen Stem Cell Conference, West Hartford, CT, August 2-3, 2018. 6th Annual NextGen Stem Cell Conference, West Hartford, CT, August 2-3, 2018. , Aug-2018

Articles in Peer-reviewed Journals

Low EK, Brudvik E, Kuhlman B, Wilson PF, Almeida-Porada G, Porada CD. "Microgravity impairs DNA damage repair in human hematopoietic stem/progenitor cells and inhibits their differentiation into dendritic cells." Stem Cells Dev. 2018 Sep 15;27(18):1257-67. Epub 2018 Jul 16. [https://](https://pubmed.ncbi.nlm.nih.gov/29901426/); PubMed PMID: 29901426, Sep-2018

Articles in Peer-reviewed Journals

Almeida-Porada G, Rodman C, Kuhlman B, Brudvik E, Moon J, George S, Guida P, Sajuthi SP, Langefeld CD, Walker SJ, Wilson PF, Porada CD. "Exposure of the bone marrow microenvironment to simulated solar and galactic cosmic radiation induces biological bystander effects on human hematopoiesis." *Stem Cells Dev.* 2018 Sep 15;27(18):1237-56. Epub 2018 Apr 26. <https://>; PubMed [PMID: 29698131](https://pubmed.ncbi.nlm.nih.gov/29698131/), Sep-2018