Fiscal Year:	FY 2022	Task Last Updated:	FY 12/16/2021
PI Name:	McGee-Lawrence, Meghan Ph.D.		
Project Title:	Osteocyte Plasma Membrane Disruptions in Sk	celetal Adaptation to Loading and	d Unloading
Division Name:	Space Biology		
Program/Discipline:			
Program/Discipline Element/Subdiscipline:			
Joint Agency Name:		TechPort:	No
Human Research Program Elements:	None		
Human Research Program Risks:	None		
Space Biology Element:	(1) Animal Biology: Vertebrate		
Space Biology Cross-Element Discipline:	(1) Musculoskeletal Biology		
Space Biology Special Category:	None		
PI Email:	mmcgeelawrence@augusta.edu	Fax:	FY
PI Organization Type:	UNIVERSITY	Phone:	706-446-0128
Organization Name:	Augusta University Research Institute, Inc.		
PI Address 1:	Department of Cell Biology and Anatomy		
PI Address 2:	1460 Laney Walker Blvd, CB1101		
PI Web Page:			
City:	Augusta	State:	GA
Zip Code:	30912	Congressional District:	12
Comments:			
Project Type:	Ground,New Investigation	Solicitation / Funding Source:	2018 Space Biology (ROSBio) NNH18ZTT001N-FG2. App D: Flight and Ground Space Biology Research
Start Date:	12/01/2020	End Date:	11/30/2023
No. of Post Docs:	1	No. of PhD Degrees:	1
No. of PhD Candidates:	4	No. of Master' Degrees:	0
No. of Master's Candidates:	0	No. of Bachelor's Degrees:	2
No. of Bachelor's Candidates:	2	Monitoring Center:	NASA ARC
Contact Monitor:	Griko, Yuri	Contact Phone:	650-604-0519
Contact Email:	Yuri.V.Griko@nasa.gov		
Flight Program:			
Flight Assignment:			
Key Personnel Changes/Previous PI:			
COI Name (Institution):	Hamrick, Mark Ph.D. (Augusta University Research Institute, Inc.) Johnson, Maribeth M.S. (Augusta University Research Institute, Inc.)		
Grant/Contract No.:	80NSSC21K0274		
Performance Goal No.:			
Performance Goal Text:			

Task Progress:Of the mouse is subjected to disuse, but the other remains mechanically loaded – allowing us to compare the effects of unloading within each animal, minimizing the number of animals needed for study and controlling for variability between mice. However, before embarking on our full studies, we first examined whether two weeks of immobilization, three weeks in tail, led to a more robust and reproducible loss of both muscle and bone mass. than our originally proposed approach. Specifically, three weeks of immobilization significantly decreased muscle mass, cortical bone area, and measurements of bone strength in the immobilized as compared to loaded limbs, whereas these trends were considerably weaker in the mice subjected to only two weeks of disuse. Therefore, we revised our experimental plans to focus on a 3 week immobilization model for all experiments moving forward, to ensure rigor and repeatability in our experiments.We are also interested in understanding what signals are specifically produced in mechanically loaded bone cells (osteocytes) that develop PMD (PMD+) as compared to cells that are loaded but do not develop PMD (PMD-). This will belp us test and establish the importance of PMD in bone's sacpation to changes in loading. Over the last year, we have developed methods to mechanically load the osteocytes, sort them based on whether they developed a PMD during loading, and then analyze the molecular signature (gene expression trends) in the PMD+ as compared to PMD-cells. These studies are still ongoing, but preliminary results suggest that the PMD+ cells are critical of initiating the earliest responses to application of a mechanical load.These studies developed the to be development of PMD with loading (by knocking out a protein called Spton1, which is involved in membrane stability), and a model where we have slowed the rate of PMD repair in osteocytes (by knockout of a protein called Spton1, which is involved in membrane st	Task Description:	The skeleton's ability to adapt to mechanical loading is crucial for bone health, as exercise promotes hypertrophy but disuse (such as from spaceflight) leads to bone loss. We were the first to report that small, transient plasma membrane disruptions (PMD) develop with in vitro and in vivo mechanical loading in bone osteocytes. These disruptions initiate skeletal mechanotransduction, suggesting PMD are stimuli recognized by osteocytes to regulate bone adaptation to its loading environment. Importantly, we consistently observe that ~20% of long bone osteocytes develop PMD with routine cage activity in mice, suggesting that formation of osteocyte PMD may be essential to bone's sensation of and response to normal gravitational loads. Accordingly, our central hypothesis is that osteocyte PMD formation is impaired during skeletal disuse, leading to bone loss. Our goals are to test the effects of disuse on osteocyte PMD formation, to determine whether osteocytes become sensitized to PMD formation and/or repair affect these processes. Our strategy is to test these concepts in an in vivo murine model of hindlimb unloading, as well as with in vitro osteocyte models of unloading (rotating wall vessel bioreactor) and reloading (fluid shear stress). Our goals align with the NASA Space Biology program as they target Research Topic 3 (Animal Biology Studies in support of Human Space Exploration)/ Sub-Topic AH1-E (Effects of fractional gravity provided by spaceflight centrifugation or ground microgravity/partial gravity analogs to gain insights into mechanisms of how animals sense, respond, and dapt to gravity shifts that are less than 1G) by discovering the contribution of osteocyte PMD formation (and hypothesized impairment during disuse) to the skeleton's adaptation to its loading environment. This project will yield a new understanding of how complex organisms adapt to the space environment, using a ground-based analog for disuse from spaceflight; we anticipate that derived data will advance strategies for skeletal mainte
Research Impact/Earth Benefits: astromatis to risk of skeletal fractures, also accion frequently on Earth in aginetis with spinal cored injures, batemis subjected to chronic/long-term bet rest, and in other cases of long-term decreased mobility. Furthermore, it is well understood that mechanical loading of the skeleton through physical exercise is beneficial for bone health across a wide spectrum of human patients, but three custinas behind how bone senses and responds to changes in mechanical loading, and exploring ways to alter the skeletal response to agiven level of mechanical loading (or withdrawal of loading), may lead to therapeutic interventions for improving bone health and reducing fracture risk. Despite this being the first year of this grant, we have made significant progress towards accomplishing the Specific Aims of the proposal. Major accomplishing the effects of reading increased cell membrane fragital volume of adaptation to loading, and testing the effects of reading increased cell membrane fragital volume of machine and increased cell membrane fragital volume on bone adaptation to loading, and testing the effects of creating delayed cell membrane fragital volume to loading, and testing the effects of oterating that mice would lose muscle and booe muscl. (beginning to discuss, that to works single hindlinb immobilization model has the advantage that one limb of the mouse is subjected to obtase. We initially proposed a two weeks single hindlinb immobilization model has the advantage that one limb of the mouse is subjected to obtase. We were ensured advantation to loading, we first camined whether two weeks of immobilization mode has the advantage that one limb of the mouse is subjected to obtase. We were ensured whyte weaks of immobilization mode has the advantage that noe limb of the mouse is subjected to obtase. The theorem main mechanical boading weeks of immobilization mode has the advantage that one limb of the	Rationale for HRP Directed Researc	h:
 Aims of the proposal. Major accomplishments include: establishment of a more robust disusc-induced bone model, beginning to define the molecular signature of mechanically loadd mechano-sensing bone cells (stocoytes) with plasma membrane disruptions (PMD), testing the effects of erating increased cell membrane fragility on bone adaptation to loading, and testing the effects of enhancing cell membrane repair on bone adaptation to loading. These are detailed below. We initially proposed a two week single hindlinb immobilization model to simulate disuse-induced loss of bone and muscle, as we had collected pilot data showing that mice would lose muscle and bone mass in this timeframe, and also show decreased evidence of osteocyte PMD in this window. This immobilization model has the advantage that one limb of the mouse is abliceted to disuse, but the other remains mechanically loadd – allowing us to compare the effects of unloading within each animal, minimizing the number of animals needed for study and controlling for variability between mice. However, before embarking on our full studies, we first examined whether two weeks of immobilization, three weeks in total, led to a more robust and reproducible loss of both muscle and bone mass than our originally proposed approach. Specifically, three weeks of immobilization significantly decreased muscle mass, cortical bone thickness, cortical bone area, and measurements of bone strength in the immobilized as compare to loaded limbs, whereas these trends were considerably weaker boins extended for only two weeks of disuse. Therefore, we revised our experiments. We are also interested in understanding what signals are specifically produced in mechanically loaded bone cells (osteocytes) that develop PMD (PMD+) as compared to effect) and repatability in our experiments. Task Progress: Task Progress: 	Research Impact/Earth Benefits:	astronauts to risk of skeletal fractures, also occurs frequently on Earth in patients with spinal cord injuries, patients subjected to chronic/long-term bed rest, and in other cases of long-term decreased mobility. Furthermore, it is well understood that mechanical loading of the skeleton through physical exercise is beneficial for bone health across a wide spectrum of human patients, but there exists a substantial proportion of the population who cannot undertake regular vigorous exercise for a variety of reasons, including underlying health conditions, time constraints, or financial concerns. Therefore, understanding the fundamental mechanisms behind how bone senses and responds to changes in mechanical loading, and exploring ways to alter the skeletal response to a given level of mechanical loading (or
preliminary results suggest that this strategy can rescue the defects caused by impaired PMD repair identified in our genetic mouse models. While we have encountered some difficulty in personnel/staff recruitment for this project due to the ongoing pandemic,	Task Progress:	Arise of the proposal. Major accomplishments include: establishment of a more robust disuse-induced bone model, beginning to define the molecular signature of mechanically loaded mechano-sensing bone cells (osteocytes) with plasma membrane disruptions (PMD), testing the effects of creating increased cell membrane fragility on bone adaptation to loading, testing the effects of creating delayed cell membrane repair on bone adaptation to loading, and testing the effects of enhancing cell membrane repair on bone adaptation to loading. These are detailed below. We initially proposed a two week single hindlimb immobilization model to stimulate disuse-induced loss of bone and subcet, as we had collected pilot data showing that mice would lose muscle and bone mass in this timeframe, and also show decreased evidence of osteocyte PMD in this window. This immobilization model has the advantage that one limb of the mouse is subjected to disuse, but the other remains mechanically loaded – allowing us to compare the effects of unloading within each animal, minimizing the number of animals needed for study and controlling for variability between mice. However, before embarking on our full studies, we first examined whether two weeks of immobilization, three weeks in total, led to a more robust and reproducible loss of both muscle and bone mass than our originally proposed approach. Specifically, three weeks of immobilization model for all experiments moving forward, to ensure rigor and repeatability in our experiments.

	we are happy to report that we have involved four PhD students, four medical students, two undergraduate students, and a high school student in completion of our funded experiments over the past year. One PhD student successfully defended her PhD and graduated earlier this year, our two undergraduate students received their Bachelor's Degrees, and all of the students involved have received authorship on either journal manuscripts or conference abstracts stemming from their contributions. Therefore, this grant is supporting the career development of the next generation of scientists.
Bibliography Type:	Description: (Last Updated: 10/24/2024)
Abstracts for Journals and Proceedings	Tuladhar A, McGee WA, Yu K, Hamrick MW, McGee-Lawrence ME. "Prkd1 is critical for repair of plasma membrane disruptions (PMD) in osteocytes." ASBMR 2021 Annual Meeting, San Diego, CA, and Virtual, October 1-4, 2021. Abstracts. ASBMR 2021 Annual Meeting, San Diego, CA, and Virtual, October 1-4, 2021. Plenary Poster, Abstract ID #A21023817. , Oct-2021
Abstracts for Journals and Proceedings	Hagan M, Piedra V, Yu K, Roberts R, Dorn J, Balayan V, Cooley M, Hamrick MW, McGee-Lawrence ME. "Sptbn1 Deficiency Blunts Adaptation In Vivo and Alters Osteocyte Plasma Membrane Dynamics And Calcium Wave Propagation In Vitro Following Formation of Plasma Membrane Disruptions (PMD)." 2021 Orthopedic Research Society, Virtual, February 12-16, 2021. Abstracts. 2021 Orthopedic Research Society, Virtual, February 12-16, 2021. ORS Annual Meeting Oral Presentation #0358. , Feb-2021
Abstracts for Journals and Proceedings	 Tuladhar A, Hagan M, Yu K, Awad M, Parker E, Hamrick MW, McGee-Lawrence ME. "Disuse from Immobilization Decreases Osteocyte Plasma Membrane Disruptions (PMD) and Causes Cortical Bone Loss." 2021 Orthopedic Research Society, Virtual, February 12-16, 2021. 2021 Orthopedic Research Society, Virtual, February 12-16, 2021.ORS Annual Meeting Poster #0531. , Feb-2021
Articles in Peer-reviewed Journals	Hagan ML, Balayan V, McGee-Lawrence ME. "Plasma membrane disruption (PMD) formation and repair in mechanosensitive tissues." Bone. 2021 Aug;149:115970. <u>https://doi.org/10.1016/j.bone.2021.115970</u> ; <u>PMID:</u> 33892174; <u>PMCID: PMC8217198</u> , Aug-2021