Fiscal Year:	FY 2009	Task Last Updated:	FY 05/18/2009		
PI Name:	Hogan, Harry Ph.D.				
Project Title:	Contributors to Long-Term Recovery of Bone Strength following Exposure to Microgravity				
Division Name:	Human Research				
Program/Discipline:	HUMAN RESEARCH				
Program/Discipline Element/Subdiscipline:	HUMAN RESEARCHBiomedical countermeasures				
Joint Agency Name:		TechPort:	No		
Human Research Program Elements:	(1) HHC :Human Health Countermeasures				
Human Research Program Risks:	 (1) Bone Fracture: Risk of Bone Fracture due to Spaceflight-induced Changes to Bone (2) Osteo: Risk Of Early Onset Osteoporosis Due To Spaceflight 				
Space Biology Element:	None				
Space Biology Cross-Element Discipline:	None				
Space Biology Special Category:	None				
PI Email:	hhogan@tamu.edu	Fax:	FY		
PI Organization Type:	UNIVERSITY	Phone:	979-845-1538		
Organization Name:	Texas A&M University				
PI Address 1:	Dept. of Mechanical Engineering				
PI Address 2:	TAMU 3123				
PI Web Page:					
City:	College Station	State:	TX		
Zip Code:	77843-3123	Congressional District:	17		
Comments:					
Project Type:	Ground	Solicitation / Funding Source:	2007 Crew Health NNJ07ZSA002N		
Start Date:	05/20/2008	End Date:	05/19/2011		
No. of Post Docs:	0	No. of PhD Degrees:	0		
No. of PhD Candidates:	0	No. of Master' Degrees:	0		
No. of Master's Candidates:	2	No. of Bachelor's Degrees:	1		
No. of Bachelor's Candidates:	3	Monitoring Center:	NASA JSC		
Contact Monitor:	Meck, J@n	Contact Phone:	281-244-5405		
Contact Email:	janice.v.meck@nasa.gov				
Flight Program:					
Flight Assignment:					
Key Personnel Changes/Previous PI:					
COI Name (Institution):	Bloomfield, Susan (Texas A&M University Martinez, Daniel (University of Houston	ity))			
Grant/Contract No.:	NNX08AQ35G				
Performance Goal No.:					
Performance Goal Text:					
	The project uses the adult male hindlimb unloaded (HU) animal model with three specific aims and associated experiments. The first aim addresses the observed "discordant recovery dynamic" reported for astronaut data (Lang et al., JBMR 21:1224, 2006) and will characterize bone mass, bone mineral density (BMD), and bone strength relationships after HU and during various periods of recovery. Additional outcome measures include bone biochemistry and gene expression. A major emphasis is to compare detailed dynamics between the animal model and astronaut data. The animal model also permits direct comparison of calculated/estimated bone strengths with measured strengths. The second aim examines multiple mission scenarios and will use HU, recovery for a period, and then a second HU exposure. The third aim will also follow the two-exposure procotol but with exercise (both aerobic and resistive) added during the recovery period.				

The cross-cutting area, or element, of the Bioastronautics Critical Path Roadmap (CRP) that this research project addresses is Human Health & Countermeasures (HHC). The specific health risk is the Risk of Accelerated Osteoporosis as identified in the Bioastronautics Roadmap (Risk No. 1, Bone Loss, p. 19 of NASA/SP–2004–6113) and the Human Research Program (HRP) Integrated Research Plan (Risk 14.0). The Gaps addressed, as defined in the HRP-IRP, are: B1 (Is bone strength completely recovered with recovery of BMD); B10 (Time-course of bone degradation during missions)

The 2007 NASA Research Announcement (NNJ07ZSA002N) to which the proposal for this project responded included the following specific solicitation wording for Gap B1: "There are preliminary indications that overall bone quality/strength does not recover at the same rate that bone mineral density recovers after spaceflight. It is not known if there is a long term health effect related to this discordant recovery dynamic." {emphasis added} Research proposals are solicited that directly address this relationship. The specific topic solicited is: Novel research that defines the precise relationship between long term recovery of bone mineral density and bone strength/quality, including the effects of multiple spaceflights." {emphasis added}

The research products to be generated from this project will mainly take the form of new knowledge about bone recovery from simulated microgravity and the response to a second exposure of simulated microgravity (following recovery from the first). In particular, insight will be gained into the interrelationships between three different, but crucial, types of bone parameters: bone mass, bone mineral density (BMD), and bone quality. Bone mass is characterized by size, shape, geometry, and bone mineral content (BMC) outcomes. Bone mineral density (BMD) has its typical meaning of the amount of mineral per unit volume, and it is thus a characteristic of bone properties at the tissue level (i.e., normalized for amount). The most basic and fundamental measure of bone quality is bone strength, and this will be measured directly with comprehensive mechanical testing of bone from multiple sites following each experiment. We will also quantify pertinent properties of the organic matrix (the unmineralized portion of bone), which is another important contributor to bone quality and forms the scaffold on which mineral nucleation occurs. Specifically, collagen content, gene expression, and cross-link maturity will be assessed. Very few of these outcome measures are available from human studies (bed-rest or flight-based). The results will also quantify the risks of previous exposure to microgravity, plus recovery, on a second exposure to microgravity.

Three sets of experiments will be conducted to address three specific aims. In all cases, adult male Sprague-Dawley rats (6-mos.-old) will be used, and the period of initial HU will be 28 days. Recovery will be characterized in two ways: (a) by comparing to age-matched, ambulatory cage control animals (no HU, but same age); and (b) by comparing to values at the end of the initial 28 days of HU (day 0 of recovery). The major outcome variables to be examined are bone mass (size, geometry, BMC), bone mineral density (total, cortical, cancellous BMD), and bone quality (strength, plus measures of tissue-level organic matrix). Tissue-level organic matrix assays will quantify collagen content, cross-link maturity, and gene expression. These will be assessed for: (i) cortical bone in the mid-diaphysis (tibia and femur); (ii) mixed cortical and cancellous bone in the metaphysis (proximal tibia and distal femur); (iii) mixed cortical and cancellous bone, both separately and combined (integrally).

The main goal of Experiment 1 is to determine the extent of loss and the time course of recovery in bone outcomes following an initial period of HU (28 days). Bone mass and density outcome variables are derived from pQCT (peripherel quantitative computed tomography) scanning, which is done both in vivo (longitudinally) and ex vivo (at each time endpoint). In vivo scans are limited to the tibia midshaft and proximal metaphysis, but ex vivo scans will be made for a full slate of anatomic sites to provide an even more thorough approach. Specifically, ex vivo pQCT results will be generated at the following locations:

Task Description:

- * Left Tibia ~ midshaft (primarily cortical bone)
- ~ proximal metaphysis (below knee, mixed cortical & cancellous)
- * Left Femur ~ midshaft (primarily cortical bone)
- ~ distal metaphysis (above knee, mixed cortical & cancellous)
- ~ femoral neck (mixed cortical & cancellous)

A distinct advantage of animal studies is that bone strength can be measured directly using harvested bones and machined specimens. The specific sites, type of test, and bone tissue are: * Left Tibia ~ 3-point bending of midshaft (primarily cortical bone)

- ~ RPC or ICC of proximal metaphysis (primarily cancellous bone)
- * Left Femur ~ 3-point bending of midshaft (primarily cortical bone)
- ~ RPC or ICC of proximal metaphysis (primarily cancellous bone)
- ~ femoral neck (mixed cortical & cancellous bone)
- * Right Femur ~ femoral neck (mixed cortical & cancellous bone)

Bone quality and underlying mechanisms will be studied and characterized through biochemical assays and gene expression quantification. This work will be done by Dr. Martinez at the University of Houston. The properties of the organic matrix will be assessed using remnants from mechanical testing specimens. The following outcome measures are planned:

* Collagen Concentration ~ hydroxyproline (Hyp) by HPLC

* Collage Cross-Links ~ non-reducible hydroxylysylpyridinoline (HP) by HPLC Six genes of interest have been identified for characterization using qRT-PCR (quantitative Reverse Transcription Polymerase Chain Reaction). These genes are:

- * Colla2 and Col3a1 ~ Extra Cellular Matrix (ECM) collagen genes
- * MMP2 or 13 ~ ECM resorption marker genes

* (Osteocal	lcin ~	mineral	homeostasis	and	osteogenic	marker
-----	----------	--------	---------	-------------	-----	------------	--------

* Runx2 and Osterix ~ mechanical loading/unloading transcription factors

The aim of Experiment 2 is to determine the precise relationships between bone mass, BMD, and bone strength after a second exposure to 28 days of HU, following an initial 28 days of HU plus a recovery period. Two recovery periods interposed between HU exposures will be examined, 28 and 56 days. The recovery periods may be modified, however, based upon results from Experiment 1. Experiment 3 will also follow the two-exposure procool but with exercise (both aerobic and resistive) added during the recovery period. Data will be analyzed to characterize and compare the effects of resistance training and treadmill running during recovery from 28 days of HU on the relationships between bone mass, BMD, and bone strength.

Bone losses observed in hindlimb unloaded adult rats parallel those in humans subjected to long-duration ISS or Mir flights; hence, this ground-based animal model is highly relevant to human astronauts. The new knowledge gained from the proposed studies will provide a better understanding of the factors affecting long term astronaut health in response to periods of exposure to microgravity. The results will provide direct, quantitative, and objective evidence for better defining the risks of space travel on long term crew member health. The results will also help define which factors are most critical to monitor in assessing recovery of bone health following single or multiple missions. From the last set of experiments, exercise countermeasure efficacy will be better understood by comparing resistance training and treadmill running when used during recovery from an initial exposure to microgravity. Results will characterize the relative effectiveness of these two types of exercise in reducing the risks of initial exposure on a second exposure to microgravity.

Rationale for HRP Directed Research:

Research Impact/Earth Benefits:	Results from this project will provide fundamental understanding of the way bone responds to mechanical unloading and how it recovers when mechanical loads are restored. Insights gained should be applicable to the clinically relevant case of aging adults with reduced activity levels. In addition, many of the same basic mechanisms overlap considerably with the broader health problem of osteoporosis and increased fracture risk in aging humans. It is widely known that bone mineral density (BMD) is not an accurate predictor of fracture incidence despite its wide use as a screening tool for osteoporosis. The findings of the research being conducted in this project will help define the relationships between BMD and other important factors, such as bone mineral content (BMC, i.e., bone mass), bone tissue quality, and most importantly bone strength. Further, the project will identify which anatomic sites in the rat provide the closest correspondence to bone loss and recovery characteristics in humans (astronauts in this case). These results should bolster the utility and robustness of rodent animal models and linking their findings to clinical cases. Finally, the project will compare and contrast the effects of aerobic and resistive exercise in restoring skeletal integrity during recovery from mechanical unloading. This information should be directly applicable to corresponding efforts aimed at using exercise to combat age-related losses from osteoporosis or related pathologies.
Task Progress:	This is a new project in its first year, so the extent of progress is limited. As the date approaches for the end of year 1 (May 19, 2009), the animal procedures for Experiment 1 are still in progress. As mentioned previously, the experiment is being conducted in 3 cohorts, and complete in vivo data are available for cohort 1 only. The trends revealed are quite encouraging, however. Specifically, plots of BMC and BMD match results for astronauts and previous HU rat studies by once again showing that BMC recovers fastest and vBMD recovers less rapidly. Ex vivo pQCT scanning is underway and will provide more detailed and thorough density results and related parameters for the wide variety of anatomic locations being studied. A notable achievement during the first year of the project was development of a new technique for mechanical testing of cancellous bone tissue from rat bones such as the femur and tibia. The proximal metaphysis of the tibia (below the knee) and the distal methapysis of the femur (above the knee) both typically contain significant proportions of both cortical and cancellous bone tissue. Our lab has previously developed a method for testing cancellous bone in specimens from these anatomic locations. The technique is called Reduced Platen Compression (RPC) testing and involves compression testing of specimens that contain both cortical and cancellous bone. This method has proved useful in quantifying changes in properties of cancellous bone, but the presence of the cortical shell as part of the speciment inherently gives rise to some degree of load sharing and a stiffer response than would be obtained with totally isolated cancellous bone specimens. Determining mechanical testing. Testing isolated samples would avoid the artifacts in RPC testing provides an effective way to do this, an effort was undertaken during year 1 to develop a way to create totally isolated cancellous bone specimens for mechanical testing. Testing isolated samples would avoid the artifacts in RPC testing due to the presen
Bibliography Type:	used for RPC testing. Next, a special fixture was designed and built to hold the slice piece for proper interfacing with a diamond wire saw (Well Diamond Wire Saws, Norcross, GA, USA). With proper placement and slow rotation of the coring fixture system, small cylindrical specimens were cut. The specimens were 2.3mm in diameter and 1.5 mm in height. Specimens were successfully created, and subsequently tested, for a bones having a wide range of densities. It was particularly encouraging that viable specimens were obtained from even the lowest density bones. Description: (Last Updated: 01/11/2021)